

# Allurion

PROGRAM



## A Comprehensive Approach to Weight Loss

- ▶ FDA approved and clinically proven to support patients to lose 2.2x more weight than with diet alone.<sup>1</sup>
- ▶ No surgery, no endoscopy<sup>2</sup>, no anesthesia.
- ▶ Placed in a brief, 15-minute visit to the doctor.
- ▶ Experience improvements in quality of life and more control over your diet.

▶ ALLURION.COM



The Allurion Gastric Balloon System (AGBS) is indicated to promote short-term limited weight loss in adult individuals with obesity between the ages of 22 years and 65 years with a body mass index (BMI)  $\geq 30$  kg/m<sup>2</sup> and  $\leq 40$  kg/m<sup>2</sup> who have had at least one unsuccessful attempt at a weight loss program. The residence time for each AGBS is variable with an average observed residence time of 15.3 weeks. The AGBS is to be used in conjunction with a moderate intensity lifestyle modification therapy program. The AGBS consists of up to two Allurion Balloons placed during a 10-month period.

1. The AUDACITY Study. 48-week LOCF results in mITT population. Data on File (2025)

2. In rare cases, the Allurion Balloon may be vomited at end of residence require endoscopic or surgical intervention for removal.

# Allurion

PROGRAM



## A Comprehensive Approach to Weight Loss

- ▶ FDA approved and clinically proven to support patients to lose 2.2x more weight than with diet alone.<sup>1</sup>
- ▶ No surgery, no endoscopy<sup>2</sup>, no anesthesia.
- ▶ Placed in a brief, 15-minute visit to the doctor.
- ▶ Experience improvements in quality of life and more control over your diet.

▶ ALLURION.COM



The Allurion Gastric Balloon System (AGBS) is indicated to promote short-term limited weight loss in adult individuals with obesity between the ages of 22 years and 65 years with a body mass index (BMI)  $\geq 30$  kg/m<sup>2</sup> and  $\leq 40$  kg/m<sup>2</sup> who have had at least one unsuccessful attempt at a weight loss program. The residence time for each AGBS is variable with an average observed residence time of 15.3 weeks. The AGBS is to be used in conjunction with a moderate intensity lifestyle modification therapy program. The AGBS consists of up to two Allurion Balloons placed during a 10-month period.

1. The AUDACITY Study. 48-week LOCF results in mITT population. Data on File (2025)

2. In rare cases, the Allurion Balloon may be vomited at end of residence require endoscopic or surgical intervention for removal.